

A Note on Quadrangular Embedding of Abelian Cayley Graphs

J.E. STRAPASSON^{1*}, S.I.R. COSTA² and M. MUNIZ³

Received on January 03, 2016 / Accepted on October 21, 2016

ABSTRACT. The genus graphs have been studied by many authors, but just a few results concerning in special cases: Planar, Toroidal, Complete, Bipartite and Cartesian Product of Bipartite. We present here a general lower bound for the genus of an abelian Cayley graph and construct a family of circulant graphs which reach this bound.

Keywords: Abelian Cayley Graphs, Genus of a graph, Flat torus, Tessellations.

1 INTRODUCTION

The genus of a graph, defined as the minimum genus of a 2-dimensional surface⁴ on which this graph can be embedded without crossings ([7, 21]), is well known as being an important measure of the graph complexity and it is related to other invariants.

A circulant graph, $C_n(a_1, \dots, a_k)$, is an homogeneous graph which can be represented (with crossings) by n vertices ($\{v_0, \dots, v_{n-1}\}$) on a circle, with two vertices being connected if and only if there is a jump of a_i from one to the other, $\forall i = 1, \dots, k$, where a jump is an edge between v_j and $v_{\text{mod}(j \pm a_i, n)}$ (Figure 1). A circulant graph is a particular case of an abelian Cayley graph. Different aspects of circulant graphs have been studied lately, either theoretically or through their applications in telecommunication networks and distributed computation [10, 12, 11, 15, 13, 9].

Concerning specifically to the genus of circulant graphs few results are known up to now. We quote [3] for a small class of toroidal (genus one) circulant graphs, [9] which establish a complete classification of planar circulant graphs, [5] which establish a complete classification of minimum genus 1 and 2 for circulant graphs, and the cases where the circulant graph is either complete or a bipartite complete graph ([4, 8, 17, 18, 20]).

*Corresponding author: João Eloi Strapasson.

¹Faculdade de Ciências Aplicadas, Unicamp – Universidade de Campinas, 13484-350 Limeira, SP, Brasil.
E-mail: joao.strapasson@fca.unicamp.br

²Departamento de Matemática, Unicamp – Universidade Estadual de Campinas, 13083-970 Campinas, SP, Brasil.
E-mail: sueli@ime.unicamp.br

³Departamento de Matemática, UFPR – Universidade Federal do Paraná, 81531-980 Curitiba, PR, Brasil.
E-mail: marcelomsa@ufpr.br

⁴The genus of a connected, orientable surface is an integer representing the maximum number of cuttings along non-intersecting closed simple curves without rendering the resultant manifold disconnected. It is equal to the number of handles on it.

In [6] the authors show how any circulant graph can be viewed as a quotient of lattices and obtain as consequences that: i) for $k = 2$, any circulant graph must be either genus one or zero (planar graph) and ii) for $k = 3$, there are circulant graphs of arbitrarily high genus.

We derive a general lower bound for the genus of abelian Cayley graph $C_n(a_1, \dots, a_k)$ as $\frac{(k - 2)n + 4}{4}$ (Proposition 1), and construct a family of abelian Cayley graphs which reach this bound (Corollary 4).

This note is organized as follows. In Section 2 we introduce concepts and previous results concerning circulant graphs, abelian Cayley graphs and genus.

In Section 3 we derive a lower bound for the genus of an n -circulant graph of order $2k$ (Proposition 1) and construct families of graphs reaching this bound for arbitrarily k (Corollary 4).

2 NOTATION AND PREVIOUS RESULTS

In this section we recall concepts and results used in this paper concerning circulant graphs. We also fix the notations which will be followed later on.

Let $G = (\{e = g_1, \dots, g_n\}, +)$ be a finite abelian group. Given a subset $S = \{a_1, \dots, a_k\}$ of G , the associated Cayley graph (G, S) is an undirected graph whose vertices are the elements of G , and where two vertices g_i and g_j are connected if and only if $g_i - g_j = \pm a_l$ for some $a_l \in S$. We remark that (G, S) is connected if and only if S generates G as a group, and that this graph is $2k$ -regular if $a_i + a_i \neq 0, \forall i = 1, 2, \dots, k$, and $(2k - l)$ -regular otherwise, where l is a number of a_i such that $a_i + a_i = 0$.

A circulant graph $C_n(a_1, \dots, a_k)$ with n vertices v_0, \dots, v_{n-1} and jumps $a_1, \dots, a_k, 0 < a_j \leq \lfloor n/2 \rfloor, a_i \neq a_j$, is an undirected graph such that each vertex $v_j, 0 \leq j \leq n - 1$, is adjacent to all the vertices $v_{j \pm a_i \pmod n}$, for $1 \leq i \leq k$. A circulant graph is homogeneous: any vertex has the same degree (number of incident edges), with is $2k$ except when $a_j = \frac{n}{2}$ for some j , when the degree is $2k - 1$, a circulant graph is a particular case of abelian Cayley graph $(G = \mathbb{Z}_n, S = \{a_1, \dots, a_k\})$.

The n -cyclic graph and the complete graph of n vertices are examples of circulant graphs denoted by $C_n(1)$ and $C_n(1, \dots, \lfloor n/2 \rfloor)$, respectively. Figure 1 shows on the left the standard picture of the circulant graph $C_{13}(1, 6)$.




Figure 1: The circulant graph $C_{13}(1, 6)$ represented in the standard form (left) and on a 2-dimensional flat torus (right).

In what follows we write $(a_1, \dots, a_k) = (\tilde{a}_1, \dots, \tilde{a}_k) \pmod n$ to indicate that for each i , there is j such that $a_i = \pm \tilde{a}_j \pmod n$. Two circulant graphs, $C_n(a_1, \dots, a_k)$ and $C_n(\tilde{a}_1, \dots, \tilde{a}_k)$ are said to satisfy the *Ádám's relation* if there is r , with $\gcd(r, n) = 1$, such that

$$(a_1, \dots, a_k) = r(\tilde{a}_1, \dots, \tilde{a}_k) \pmod n \quad (1)$$

An important result concerning circulant graphs isomorphisms is that circulant graphs satisfying the Ádám's relation are isomorphic ([1]). The reciprocal of this statement was also conjectured by Ádám. It is false for general circulant graphs but it is true in special cases such as $k = 2$ or $n = p$ or $n = pq$ (p and q prime) (see [11, 2]). In this paper we will not distinguish between isomorphic graphs.

Without loss of generality we will always consider $a_1 < \dots < a_k \leq n/2$ for a circulant graph $C_n(a_1, \dots, a_k)$.

A circulant graph $C_n(a_1, \dots, a_k)$ is connected if, and only if, $\gcd(a_1, \dots, a_k, n) = 1$ ([3]). *In this paper we just consider connected circulant graphs.*

The genus of a graph is defined as the minimum genus, g , of a 2-dimensional orientable compact surface \mathcal{M}_g on which this graph can be embedded without crossings ([7, 21]). This number, besides being a measure of the graph complexity, is related to other invariants. Let G a graph of the genus g , defines p_g as $p_g = \left\lfloor \frac{(7 + \sqrt{48g + 1})}{2} \right\rfloor$, so: the *chromatic number* of G is $\gamma(G) = p_g$ (Heawood conjecture) and the *algebraic connectivity*⁵ of G , $\mu(G)$, satisfies $\mu(G) < p_g - 1$ for all noncomplete graphs G if $p_g(p_g - 7) = 12(g - 1)$, see [14].

A graph E is a *subdivision* of H if it is constructed from H by possibly adding new vertices on the edges of H . Finally, if there is a subdivision E of H which is a subgraph of G we say G is *supergraph* of H . From this definition follows that

$$\text{if } G \text{ is a supergraph of } H, \text{ genus}(G) \geq \text{genus}(H).$$

When a connected graph G is embedded on a surface, \mathcal{M}_g , of minimum genus g it splits the surface in regions called *faces*, each one homeomorphic to an open disc surrounded by the graph edges, giving rise to a tessellation on this surface. Denoting the number of faces, edges and vertices by f , e , and v respectively, those numbers must satisfy the well known Euler's second relation:

$$v + f - e = 2 - 2g \quad (2)$$

We quote next other known relations those numbers must satisfy ([7, 21]):

If G is a graph of genus g with $v \geq l$ such that any face in \mathcal{M}_g has at least l sides in its boundary,

$$lf \leq 2e \text{ and } g \geq \frac{l-2}{2l}e - \frac{1}{2}(v-2). \quad (3)$$

In the above expressions we have equalities if, only if, all the faces have l sides.

An upper bound for the genus of a connected graph of n vertices is given by the genus of the *complete graph*, $C_n(1, \dots, \lfloor n/2 \rfloor)$, which is $\left\lceil \frac{(n-3)(n-4)}{12} \right\rceil$. Combining the lower bound above with a minimum of three edges for each face, we can write the following inequality, for $n \geq 3$:

$$\left\lceil \frac{1}{6}e - \frac{1}{2}(n-2) \right\rceil \leq g \leq \left\lceil \frac{(n-3)(n-4)}{12} \right\rceil, \quad (4)$$

⁵The algebraic connectivity is the second-smallest eigenvalue of the Laplacian matrix of G .

where $\lceil x \rceil$ is the ceiling (smallest integer which is greater or equal to) of x .

For a circulant graph $C_n(a_1, \dots, a_k)$, $a_1 < a_2 < \dots < a_k$ we can replace e by $e = nk$ when $a_k < \frac{n}{2}$, or $e = n(2k - 1)/2$ when $a_k = \frac{n}{2}$. We can then rewrite the lower bound in last expression as $\lceil \frac{n}{6}(k - 3) + 1 \rceil$ or $\lceil \frac{n}{6}(k - 4) + 1 \rceil$, respectively.

2.1 Previous results on genus of circulant graphs and abelian Cayley graphs

- **Theorem** (Ringel, Beineke and Harary, 1965 [19, 4]). *The genus of the n -cube graph Q_n is $1 + 2^{n-3}(n - 4)$.*
- **Theorem** (Ringel, 1965 [17, 18]). *The genus of the complete bipartite graph $K_{m,n}$ is $\lceil \frac{(m-2)(n-2)}{4} \rceil$.*
 Since $K_{n,n}$ is the circulant graph $C_{2n}(1, 3, \dots, 2 \lceil \frac{n-1}{2} \rceil - 1)$, the genus of this one-parameter family is $\lceil \frac{(n-2)^2}{4} \rceil$.
- **Theorem** (White, 1970 [22]). *Let $G = C_{m_1} \square C_{m_2} \dots \square C_{m_r}$ ⁶, where C_{m_i} is even cycle, $r > 1$ and $m_i > 3$ for all i . Then the genus of G is $1 + v(r - 2)/4$, where $v = m_1 m_2 \dots m_r$.*
- **Theorem** (Pisanski, 1980 [16]) *Let G and H be connected r -regular bipartites graphs. Then the Cartesian product $G \square H$ of G and H has genus $1 + pm(r - 2)/4$ where p and m are the number of vertices of G and H , respectively.*
- **Theorem** (Heuberger, 2003 [9]). *A planar circulant graph is either the graph $C_n(1)$, or $C_n(a_1, a_2)$, where i) $a_2 = \pm 2a_1 \pmod n$ and n is even, ii) $a_2 = n/2$, and a_2 is even.*
- For $k = 2$, and general (a_1, a_2) , we have shown that circulant graphs $C_n(a_1, a_2)$ are very far from reaching the upper bound for the genus given in (4), as it was shown in [6]:

Proposition 1 ([6]). *Any circulant graph $C_n(a_1, a_2)$, $a_1 < a_2 \leq n/2$, has genus one, except for the cases of planar graphs: i) $a_2 = \pm 2a_1 \pmod n$, and n is even, ii) $a_2 = n/2$, and a_2 is even.*

- For $k = 3$ and $n \neq 2a_3$ we can assert that the genus of $C_n(a_1, a_2, a_3)$ satisfies:

$$1 \leq g \leq \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil \tag{5}$$

The genus of the complete graph $C_7(1, 2, 3)$ achieves the minimum value one (4). However, in opposition to the case $k = 2$, the genus of a circulant graph $C_n(a_1, a_2, a_3)$ can be arbitrarily high:

Proposition 2 ([6]). *There are circulant graphs $C_n(a_1, a_2, a_3)$ of arbitrarily high genus. A family of such graphs is given by: $n = (2m + 1)(2m + 2)(2m + 3)$, $m \geq 2$; $a_1 = (2m + 2)(2m + 3)$, $a_2 = (2m + 1)(2m + 2)(m + 1)$, $a_3 = (2m + 2)(2m + 3)(m + 1)$, with the correspondent genus satisfying*

$$g \geq 2m(m + 1)^2 + 1. \tag{6}$$

In the next section we deal with the more general class of abelian Cayley graphs and establish a lower bound for their genus.

⁶The Cartesian product $G_1 \square G_2$ of the graphs G_1 and G_2 is a graph such that the vertex set of $G_1 \square G_2$ is the Cartesian product of the set of vertices of G_1 with the set of vertices G_2 ($V(G_1) \times V(G_2)$) and any two vertices (u, u') and (v, v') are adjacent in $G_1 \square G_2$ if and only if either $u = v$ and u' is adjacent with v' in G_2 , or $u' = v'$ and u is adjacent with v in G_1 .

3 QUADRANGULAR EMBEDDING OF ABELIAN CAYLEY GRAPHS

In this section we consider Cayley graphs of abelian groups, a more general class of graphs of which circulant graphs form a very particular subclass, the Cayley graphs of cyclic groups. Nevertheless, an important feature of circulant graphs, their embeddings in k -dimensional tori, is shared by the whole class of Cayley graphs of abelian groups. We will see in the following that k is associated to the number of elements of the generating set of the edges of the Cayley graph. We will determine a subclass of these graphs that has quadrangular embeddings, and hence a subclass where we know the genus of each graph.

It is known that graphs that have 3-cycles may have embeddings with triangular faces, and some easy calculations establish a lower bound for the genus. In general we can also establish a lower bound that depends on the girth l of the graph. If (G, S) is a Cayley graph and there are no solutions of $a_h = \pm(a_i \pm a_j)$ for $h, i, j \in \{1, 2, \dots, k\}$ (not necessarily distinct), the girth is always 4 (a typical 4-cycle is $0, a_i, a_i + a_j, a_j, 0$), which implies at least four edges for each face.

If the graph is $2k$ -regular, then

$$g \geq \frac{l-2}{2l} a - \frac{v-2}{2} = \frac{2}{8} nk - \frac{n-2}{2} = \frac{nk - 2n + 4}{4}.$$

Hence, we get the following lemma, which establishes a lower bound for circulant graphs with no triangular faces.

Lemma 1. *The genus, g , of the circulant graph $C_n(a_1, \dots, a_k)$, such that $a_i \neq a_j + a_l, \forall i, j, l \leq k$ and $n \neq 2a_i, \forall i$ satisfy:*

$$g \geq \frac{nk - 2n + 4}{4}.$$

In what follows the (additive) subgroup of $G = \mathbb{Z}_{n_1} \times \mathbb{Z}_{n_2} \times \dots \times \mathbb{Z}_{n_l}$ generated by $a_1, \dots, a_k \in G$ is denoted by $\langle a_1, \dots, a_k \rangle$, and let $G_s = \langle a_1, a_2, \dots, a_s \rangle \triangleleft G$. Define L_s as the group order of $a_s, 1 \leq s \leq k$, where $L_1 = o(a_1)$ is the group order of a_1 , and $L_s = o(a_s + G_{s-1}) = [G_s : G_{s-1}]$ is the index of the quotient group $1 < s \leq k$.

Under the above conditions we can assert that $x \in G$ can be expressed uniquely as a linear combination, $x = m_1 a_1 + m_2 a_2 + \dots + m_k a_k$, where $0 \leq m_i < L_i$. This fact is stated in the next a lemma.

Lemma 2. *Given $x \in G$ and $G_s \triangleleft G, 1 \leq s < k$ then there exist a unique $m_i \in \mathbb{N}$ and $R_{s,x}$ such that*

$$x = m_1 a_1 + \dots + m_s a_s + R_{s,x} \text{ and } R_{s,x} = R(m_{s+1}, \dots, m_k) = m_{s+1} a_{s+1} + \dots + m_k a_k$$

with $0 \leq m_i < L_i$ for all i .

Through the Lemma 2 we can show that not only the circulant graphs [6] but any Cayley graph of an abelian group can be embedded in a k -dimensional torus. The construction of such embedding, for $k = 2$, is illustrate in Figure 2.

We consider the mapping:


$$\begin{aligned} \varphi : \quad \mathbb{Z}^k &\longrightarrow G \\ (x_1, \dots, x_k) &\longmapsto x_1 a_1 + \dots + x_k a_k. \end{aligned} \tag{7}$$

Therefore $\frac{\mathbb{Z}^k}{\ker \varphi} \simeq G$ and $\ker \varphi$ is lattice. The Cayley graph associated to G , as a quotient of lattices, is then naturally embedded in flat torus which a polytope generated by basis of this lattice, with the parallel faces identified.

To proceed in a uniform way we can use the standard Hermite basis for $\ker \varphi$, as it done for circulant graphs in [9].

We remark that Hermite basis of $\ker \varphi$, $\{U_1, \dots, U_k\}$, is given as columns of a upper triangular matrix $(b_{i,j})_{k \times k}$, where $b_{i,i} = L_i$ and $0 \leq b_{i,j} < L_i$.

In Figure 2, we consider the Cayley graph of $G = \mathbb{Z}_2 \times \mathbb{Z}_8$ and $a_1 = (1, 2)$ and $a_2 = (0, 1)$, therefore $o(a_1) = 4$ and $o(a_2 + \langle a_1 \rangle) = 4$ and Hermite basis is $\{(4, 0), (2, 4)\}$.



(a) Flat torus



(b) Torus

Figure 2: 2-embedding of Cayley graph $(\mathbb{Z}_2 \times \mathbb{Z}_8, \{(1, 2), (0, 1)\})$.

We will construct the embedding of Cayley graphs (G, S) by induction on $k = \#S$ ($\#S$ is equal to the cardinality of S). Note that $(G, S - \{a_k\})$ may be disconnected: it is well known that this graph has $d = \gcd(n, a_1, \dots, a_{k-1})$ components, where $L_k = [G = G_k : G_{k-1}] = o(a_k + G_{k-1})$, and that each component is isomorphic to the Cayley graph $(G_{k-1}, S - \{a_k\})$. Since x and y are linked by a path if and only if $x - y = m_1 a_1 + \dots + m_{k-1} a_{k-1}$ in G , it follows that x and y are in the same component if and only if $x \equiv y \in G_{k-1}$. Hence, each $0 \leq j_s < L_k$ determines a component and the numbers,

$$R_{k,m_k} + m_1 a_1 + \dots + m_{k-1} a_{k-1}$$

describe all vertices of the component of $(G, S - a_k)$ associated to m_k , where $0 \leq m_k < L_k$ and R_{k,m_k} is a fixed element of this component.

Proposition 3. Let $(G, \{a_1, \dots, a_k\})$, where $n = \#G = 2l$, and $L_i = 2l_i$, $1 \leq i \leq k$, and $l_1 > 1$ and $a_i \neq \pm(a_j \pm a_h)$, $1 \leq i, j, h < k$. Hence, the genus of G is $\frac{nk - 2n + 4}{4}$.

Proof. The proof will be done by induction on k . For $k = 2$ it is trivial (vide Figure 2). Assume that the result holds for $k - 1$. The graph $(G, \{a_1, \dots, a_{k-1}\})$ is a disconnected Cayley graph with an even number of connected components $(L_k = 2l_k)$, $H_{m_k} = R_{k,m_k} + G_{k-1}$, $0 \leq m_k < 2l_k$, where $2l_k = [G : G_{k-1}]$. Each component H_{m_k} can be embedded on a surface S_{m_k} giving to a rise tessellation where every face has 4 edges. As in Figure 3, we reverse the orientation of the components that contain the odd multiples of a_k . We wish to add tubes that, topologically, are prisms with squared bases.




Figure 3: The construction of the $C_{32}(8, 2, 3)$ embedding. Tubes are added on the two connected components of $C_{32}(8, 2)$ considered with reversal orientation.

Let $x \in H_{m_k}$, then $x = m_k a_k + m_1 a_1 + \dots + m_{k-1} a_{k-1}$ where $0 \leq m_k < 2l_k$ and $0 \leq m_i < 2l_i$ as in Lemma 2. Given $0 < j < k - 1$, we can also express

$$x = m_k a_k + (2p_j + \delta_j) a_j + (2p_{k-1} + \delta_{k-1}) a_{k-1} + g \tag{8}$$

$$x = m_k a_k + (2q_j + \delta_j) a_j + ((2q_{k-1} + 1) - \delta_{k-1}) a_{k-1} + g \quad (q_j > 0) \tag{9}$$

with $g = \sum_{\substack{i=1, \dots, k-2 \\ i \neq j}} m_i a_i$ and $\delta_j, \delta_{k-1} \in \{0, 1\}$.

Three possibilities should be considered:

- i) Each vertex of H_{m_k} , m_k even, is a vertex of a square determined by $\{P, P + a_{k-1}, P + a_{k-1} + a_j, P + a_j\}$, where P is of the form

$$P = m_k a_k + 2p_j a_j + 2p_{k-1} a_{k-1} + g, \tag{10}$$

where $0 \leq 2p_j < L_j, 2p_{k-1} < L_{k-1}$. Just as Figure 3, each such square is then connected to the square $\{P + a_k, P + a_k + a_{k-1}, P + a_k + a_{k-1} + a_j, P + a_k + a_j\}$ (which lies on $H_{m_{k+1}}$) by a prism which contains the edges $[P, P + a_k], [P + a_{k-1}, P + a_k + a_{k-1}], [P + a_{k-1} + a_j, P + a_k + a_{k-1} + a_j]$ and $[P + a_j, P + a_k + a_j]$; and then we cut out both squares. Doing this for every $j \in \{1, 2, \dots, k-2\}$ we construct a surface where each edge of the form $[x, x + a_k]$ is embedded without crossings.

- ii) Each vertex of H_{m_k} , m_k odd and $m_k \neq 2l_k - 1$, is a vertex of a square determined by $\{Q, Q + a_{k-1}, Q + a_{k-1} + a_j, Q + a_j\}$, where Q is of the form

$$Q = m_k a_k + 2q_j a_j + (2q_{k-1} + 1) a_{k-1} + g, \tag{11}$$

where $2 \leq 2q_j \leq L_j, 2q_{k-1} < L_{k-1}$. The same reasoning as above can be applied by replacing P by Q .

- iii) Each vertex, x of $H_{2l_{k-1}}$, $x + a_k \in H_0$. This case requires special care, since we need to choose a face in $H_{2l_{k-1}}$ and another in H_0 , once that some faces have been excluded. This choice depends on how $2l_k a_k$ is described in G_k , since $2l_k a_k = \tilde{m}_j a_j + \tilde{m}_{k-1} a_{k-1} + \tilde{g}$, $\tilde{g} = \sum_{\substack{i=1, \dots, k-2 \\ i \neq j}} \tilde{m}_i a_i$, choose

\tilde{Q} such that $m_j + \tilde{m}_j, m_{k-1} + \tilde{m}_{k-1}$ and m_j are not both even, and m_{k-1} is not odd. We then repeat the procedure of item (i), replacing P by \tilde{Q} .

Therefore, under restrictions considered in this proposition we always can connect the excluded squares by prisms and construct a surface which is tessellated by (G, S) , and each face of this tessellation is a square, by Lemma 1 the genus is greater than or equal to $\frac{nk - 2n + 4}{4}$, remember that equality is achieved if all faces have the same number of sides, and this concludes the proof. \square

Corollary 4. *Let $G = C_n(a_1, \dots, a_k)$, where $n = 2^r l$, $a_i = 2^{r_i} l_i$, $i = 1, \dots, k - 1$, where l, l_i, a_k odd, $0 < r_{i+1} < r_i < r$ and $a_{i+1} \neq \pm 2 a_i$, $1 \leq i < k$. Hence, the genus of G is $\frac{nk - 2n + 4}{4}$.*

The next example shows that there are more circulant graphs than the ones considered in Proposition 3 which also can be embedded giving rise to a quadrilateral tessellation.

Example 3.1. *For the graph $C_{32}(8, 2, 3, 7)$, if we consider $C_{32}(8, 2, 3)$ as in last proposition, we note that just half the faces of the tessellation of $C_{32}(8, 2)$ are excluded to add tubes. We can also exclude the other faces adding tubes to support the edges $\pm a_4$. Hence this is an embedding generating quadrilateral faces and since there are no cycles of size 3, the expression $\frac{nk - 2n + 4}{4}$ for the genus still holds.*

Figure 4 shows all the circulant graphs of 32 vertices for which the genus can be given by Proposition 2.1 (Heuberger), 1 and Corollary 4

k	$a_1 \in$	$a_2 \in$	$a_3 \in$	$a_4 \in$	g
1	I				0
2	I	$2(I - \{\pm a_1\})$			1
2	I	$4I$			1
2	I	$8I$	-	-	1
3	I	$2(I - \{\pm a_1\})$	$4(I - \{\pm 2 a_2\})$	-	9
3	I	$2(I - \{\pm a_1\})$	$8I$	-	9
3	I	$4I$	$8(I - \{\pm 2 a_2\})$	-	9
4	I	$2(I - \{\pm a_1\})$	$4(I - \{\pm 2 a_2\})$	$8(I - \{\pm 2 a_3\})$	17

Figure 4: All circulant graphs of 32 vertices satisfying Propositions 2.1 (Heuberger), 1 and 4 ($I = \{\pm 1, \pm 3, \dots, \pm 15\}$).

We note that some graphs satisfying the hypotheses of the last proposition belong to the class of graphs with given genus. For some of those graphs we could have used the results of White and Pisanski (see [22, 16]) to determine their genus namely. The particular class of Cartesian product of bipartite graphs which satisfy the proposition hypothesis. However, many of the graphs considered in the last proposition are not Cartesian product of bipartite graphs.

For example the Cayley graphs $G_1 = (\mathbb{Z}_2 \times \mathbb{Z}_8, \{(1, 2), (0, 1)\})$ and $G_2 = C_{16}(1, 4)$.

We can assert that the G_1 is not a product of de bipartite graphs, since as $G_1 = G \square H$, a regular graph, to be such a product, each graph factor needed to be also a regular graph. There are few possibilities to be considered here: $\#G = 2$ or 4 and $\#H = 8$ or 4 , respectively. That is, either i) $G = P_2$ and H is a 3-regular complete bipartite graph (hypothesis of White and Pisanski results), what is not possible since H is bipartite and therefore it should be 4-regular, or ii) $G = H = K_{2,2}$ what again cannot be true, since

$K_{2,2} \square K_{2,2} = (\mathbb{Z}_4 \times \mathbb{Z}_4, \{(1, 0), (0, 1)\})$, which has spectrum $\{8, 6, 6, 6, 6, 4, 4, 4, 4, 4, 4, 2, 2, 2, 2, 0\}$ and G_1 has spectrum $\{8, 6, 6, 4 + \sqrt{2}, 4 + \sqrt{2}, 4 + \sqrt{2}, 4 + \sqrt{2}, 4, 4, 4 - \sqrt{2}, 4 - \sqrt{2}, 4 - \sqrt{2}, 4 - \sqrt{2}, 2, 2, 0\}$.

G_2 also is not a product of bipartite graphs since it has odd size cycles (ex: 0, 1, 2, 3, 4, 0) what does not occur for bipartite graphs.

4 ACKNOWLEDGMENTS

This work was partially supported by FAPESP-Brazil 2007/56052-8, 2007/00514-3, 2011/01096-6 and 2013/25977-7 and CNPq 309561/2009-4 and 304705/2010-1.

RESUMO. O gênero de grafos têm sido estudados por muitos autores, mas existem resultados apenas para casos especiais: Planar, Toroidal, Completo, Bipartido e Produto cartesiano de Bipartidos. Apresentamos aqui um limite inferior para o gênero de um grafo de Cayley de um grupo abeliano e construímos uma família de grafos circulantes que atingem esse limitante.

Palavras-chave: Grafos de Cayley abelianos, Gênero de grafos, Toro plano, Tesselacões.

REFERENCES

- [1] A. Ádám. Research problem 2-10. *J. Combinatorial Theory*, (1967).
- [2] B. Alspach & T.D. Parsons. Isomorphism of circulant graphs and digraphs. *Discrete Math.*, **25**(2) (1979), 97–108.
- [3] F. Boesch & R. Tindell. Circulants and their connectivities. *J. Graph Theory*, **8**(4) (1984), 487–499.
- [4] L. W. Beineke & F. Harary. The Genus of the n-Cube. *Canad. J. Math.*, **17** (1965), 494–496.
- [5] M. Conder & R. Grande. On Embeddings of Circulant Graphs. *The Electronic Journal of Combinatorics*, **22**(2) (2015), 1–27.
- [6] S.I.R Costa, J.E Strapasson, M.Muniz & T.B. Carlos. Circulant graphs and tessellations on flat tori. *Linear Algebra and Appl*, **432**(1) (2010), 369–382.
- [7] J.E. Gross & T.W. Tucker. *Topological graph theory*. Dover Publications Inc., Mineola, NY (2001).
- [8] F. Harary, *Graph Theory*. Reading, MA: Addison-Wesley, (1994).
- [9] C. Heuberger. On planarity and colorability of circulant graphs. *Discrete Math.*, **268**(1-3) (2003), 153–169.
- [10] X. Lin, Y. Yang, J. Lü & X. Hao. The crossing number of $C(mk; \{1, k\})$. *Graphs Combin.*, **21**(1) (2005), 89–96.
- [11] V. Liskovets & R. Pöschel. Counting circulant graphs of prime-power order by decomposing into orbit enumeration problems. *Discrete Math.*, **214**(1-3) (2000), 173–191.
- [12] D. Matthew. On Hamilton cycle decomposition of 6-regular circulant graphs. *Graphs Combin.*, **22**(3) (2006), 331–340.
- [13] F.P. Muga II. Undirected Circulant Graphs, International Symposium on Parallel Architectures. *Algorithms and Networks*, (1994), 113–118.
- [14] J.J. Molitierno. On the algebraic connectivity of graphs as a function of genus. *Linear Algebra and its Applications*, **419**(2-3) (2006), 519–531.

- [15] M. Muzychuk. \acute{A} dám's conjecture is true in the square-free case. *J. Combin. Theory Ser. A*, **72**(1) (1995), 118–134.
- [16] T. Pisanski. Genus of Cartesian products of regular bipartite graphs. *J. Graph Theory*, **4** (1980), 31–42.
- [17] G. Ringel. Das Geschlecht des vollständiger Paaren Graphen. *Abh. Math. Sem. Univ. Hamburg*, **28** (1965), 139–150.
- [18] G. Ringel, Der vollständige paare Graph auf nichtorientierbaren Flächen. *J. Reine Angew. Math.*, **220** (1965), 88–93.
- [19] G. Ringel, Über drei kombinatorische Problem am n-dimensionalen Würfel und Wurfelgitter. *Abh. Math. Sem. Univ. Hamburg*, **20** (1965), 10–19.
- [20] G. Ringel & J.W.T. Youngs. Solution of the Heawood Map-Coloring Problem. *Proc. Nat. Acad. Sci. USA*, **60** (1968), 438–445.
- [21] R.J. Trudeau. *Introduction to graph theory*. Dover Publications Inc., New York, 1993. Corrected reprint of the 1976 original.
- [22] A.T. White. The genus of repeated cartesian products of bipartite graphs. *Trans. Amer. Math. Soc.*, **151** (1970), 393–404.